



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 5/00, A61K 45/05, 48/00, 31/40	A1	(11) International Publication Number: WO 96/29394 (43) International Publication Date: 26 September 1996 (26.09.96)
(21) International Application Number: PCT/US96/03621 (22) International Filing Date: 15 March 1996 (15.03.96) (30) Priority Data: 08/406,388 17 March 1995 (17.03.95) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). (72) Inventor: GRANGER, Gale, A.; 31562 Santa Rosa, Laguna Beach, CA 92677 (US). (74) Agent: BERLINER, Robert; Robbins, Berliner & Carson, 5th floor, 201 North Figueroa Street, Los Angeles, CA 90012-2628 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: METHOD FOR TREATING TUMORS (57) Abstract A method is provided for treatment of a mammalian patient having a tumor by administering to the patient allogenic donor lymphocytes that have been co-cultured in the presence of the patient-derived lymphocytes under conditions sufficient to alloactivate the donor lymphocytes. It is preferred that the donor lymphocytes be introduced intralesionally. This method is preferred for treatment of glioblastoma in humans.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

METHOD FOR TREATING TUMORS

BACKGROUND OF THE INVENTION

5 This application is a continuation-in-part application of U.S. patent application Serial No. 08/406,388, filed March 17, 1995.

1. *Field of the Invention*

10 This invention relates to methods for inhibiting tumor cell proliferation. More particularly this invention relates to a method for inhibiting tumor cell proliferation by enhancing the patient's ability to respond immunologically to the tumor.

15 2. *Description of Related Art*

Traditional therapies have done little to alter the outcome for patients with high-grade brain tumors, such as glioblastomas, and many other types of tumors, such as systemic melanoma, and cancers of the head and neck.

20 Patients with resectable primary tumors generally experience recurrence of the tumor within one year after surgery, chemotherapy, or radiation. Often these tumors progress rapidly, with or without further conventional therapy. Thus, there is the need to develop new modes of therapy for these deadly tumors.

25 A new family of cancer therapies developed in recent years are based on immunotherapy. In general, tumor immunotherapies take one of two approaches: 1) various techniques are employed to activate the patient's immune system to attack the tumor; or 2) the lymphoid cells of the patient are removed and activated by *in vitro* techniques to produce anti-cancer
30 activity, and the activated cells are then systemically reintroduced into the

patient. The clinical effectiveness of these various types of immunotherapy are being evaluated in patients with different types of cancers. However, one of the major problems associated with both of these types of immunotherapy is toxicity observed when immunotherapeutic agents are administered systemically. A method developed to avoid this toxicity is intralesional administration of immunotherapy, for instance by injection directly into the tumor. Intralesional administration of various forms of immunotherapy to cancer patients does not cause the toxicity seen with systemic administration of immunologic agents (M. Fletcher, *et al.*, *Lymphokine Res.* 6:45, 1987; H. Rabinowich, *et al.*, *Cancer Res.* 47:173, 1987; S.A. Rosenberg, *et al.*, *Science* 233:1318, 1989; and G. Pizz, *et al.*, *Int. J. Cancer* 34:359, 1984).

Recent studies indicated that immunization of animals with tumor cells that were genetically engineered to secrete different cytokines enhanced the induction of a therapeutic immune response. The cytokines are believed to induce a complex set of reactions including: a) increased expression of tumor antigens; b) inflammation and infiltration of the tumor with host lymphoid cells; c) induction of tumor specific immunity; and d) activation of both nonspecific and specific host anti-tumor effector mechanisms, which destroy the tumor. However, although this technique may ultimately prove useful, because it is extremely costly and time consuming, its application may be limited.

Studies in experimental animals (primarily mice) show that the chronic release of cytokines within a tumor may induce a host anti-tumor response and tumor regression. Repeated intralesional injection of cytokines such as Interleukin-2 (IL-2), Tumor Necrosis Factor (TNF) and Interferon- γ (INF- γ) has been shown to cause regression of cutaneous sarcomas (S.P. Creekmore, *et al.*, *Resident and Staff Physician* 34:23-31, 1988; P. Greenberg, *et al.*, *Basic and Tumor Immunology* (R. Herberman, ed.) p. 302, 1983; E. Grimm,

et al., *Lymphokines*, 9:279-311, 1984; G. Forni, *et al.*, *Lymphokines* 14:335-360, 1987). It has also been shown that injection into a tumor of the animal's tumor cells that have been genetically engineered to secrete cytokines such as IL-2, IL-4, TNF and Granulocyte Monocyte Colony Stimulating Factor (GM-CSF) will induce host anti-tumor immunity (E. Feron, *et al.*, *Cell* 60, 397-403, 1990; P. Galumbek, *et al.*, *Science* 254:713-716, 1991; A. Ascher, *et al.*, *J. of Immunol.* 146:3227-3234, 1991). These latter results have been obtained even in treatment of tumors that were previously thought to be non-immunogenic.

10

J.M. Redd, *et al.* (*Cancer Immunology and Immunotherapy* 34(5):349, 1992) have shown in rats that allogeneic lymphocytes sensitized against donor alloantigens can inhibit tumor formation when co-injected into the brain of a rat with 9L glioblastoma. In a separate study, both normal and alloimmune spleen cells from Wistar rats were injected into established 6-day T9 brain tumors in the Fischer rat. Intralesional injection of normal Wistar spleen cells from Wistar rats, previously immunized against Fischer alloantigens, cured the tumors in 50% of the Fischer rats. In contrast, untreated animals and non-responders died within 30 days. Survivors appeared completely normal and intracranial injections of allogeneic cells into normal rats caused no detectable change in behavior or survival over a three month period. Histopathologic examination of brains from treated tumor bearing animals revealed: a) mononuclear cell infiltration, massive tumor necrosis beginning at 2 to 4 days and total tumor destruction by 15 days; or 2) cellular infiltration, early tumor destruction and then tumor regrowth progressing to death of the animal. No damage to normal brain tissue was evident at any time in these animals. Tumor regressor animals developed systemic immunity, for they proved totally resistant to intracranial rechallenge with viable tumor. Although these results in rats are of interest, their value in reasonably predicting what would be seen in a highly unrelated species, such

30

as a human, is highly questionable in view of the considerable species diversity which exists, especially with respect to the immunological response to tumors.

5 Human glioma patients with localized and surgically accessible tumors are logical candidates for intralesional immunotherapy. Multiple Phase I studies in adult patients with gliomas have been reported employing intratumor implants of autologous peripheral blood lymphocytes activated *in vitro* with IL-2. While little clinical effect was noted, negative side effects were few
10 and occurred only when excessive levels of IL-2 were co-administered with the cells (K.S. Jacobs, *et al.*, *Cancer Res.*, 47:2101, 1986; R. Merchant, *et al.*, *Neurosurgery* 23:725, 1988). Studies of these patients revealed that survival correlated directly with the ability of implanted cells to secrete the cytokine TNF. The discovery of inhibitors for both TNF and IL-1 in the serum or
15 tumor cyst fluid, and in primary cultures of the tumors from these patients suggests the tumor cells surround themselves with agents that block the host anti-tumor response. These inhibitors may prevent cytokine activated implanted cells from remaining active in the tumor long enough to cause its destruction. This concept is supported by findings in brain tumor studies in
20 rats. When IL-2 active lymphoid cells were implanted into C6 and T9 glioma brain tumors in Wistar and Fischer rats, respectively, histopathologic examination revealed the implanted IL-2 activated lymphoid cells only remained in the tumor site for 4 to 6 days (W. Carson *et al.*, *J. of Immunotherapy* 10(2):131-140, 1991).

25

The mechanisms operative in causing tumor regression in the animals treated with allogeneic lymphoid cells include a graft vs. host and host vs. graft reaction in the tumor site. These powerful immunologic reactions may stimulate high levels of endogenous cytokine production in the tumor,
30 overcome local levels of cytokine inhibitors, and presumably stimulate

infiltration, recruitment and activation of both specific and non-specific host anti-tumor activity. The tumor regressor animals were found to be resistant to tumor rechallenge. However, it has heretofore been unknown whether any treatments based on similar methods would achieve similar results in human subjects sufficient to be considered effective in the treatment of human tumors.

Therefore, in view of the limitations of the prior art, new and better methods for treating mammalian tumors are needed. In particular, new methods of intratumor immunotherapy are needed for human cancer patients for whom regression of an individual solid tumor could prove life saving.

SUMMARY OF THE INVENTION

The present invention provides a method for inhibiting tumors in mammals. In the general practice of this invention, the tumor patient's peripheral blood mononuclear cells (PBMCs) are co-cultured *in vitro* to induce a mixed lymphocyte cell reaction with healthy lymphocytes derived from a normal donor, preferably an allogeneic donor. Preferably the normal lymphocytes are from a donor unrelated to the tumor patient, and preferably the normal lymphocytes are obtained by leukapheresis from whole blood. During co-culture the allogeneic donor lymphocytes are specifically activated against the patient's alloantigens. The mixture of live alloactivated donor and patient lymphocytes produced by mixed lymphocyte culture are referred to herein as "MLCs." The MLC activated cells produce a mixture of cytokines which have been shown to induce a primary immune response *in vitro*. In the treatment of glioblastoma, for instance, the patient is a human and the alloactivated lymphocytes are surgically implanted into the patient's brain at the primary tumor site, optionally together with the patient's PBMCs from the co-culture as a MLC mixture, to induce the patient's immune system to attack autologous tumor cells. In the treatment of other types of tumors, the MLCs are injected into a tumor site, site of metastases, or body cavity, such as the peritoneum. Optionally, the MLCs are administered peripherally, such as at a non-primary tumor site, in the treatment of tumors other than glioblastoma.

In one embodiment of the invention, the alloactivated donor lymphocytes obtained by co-cultivation with patient-derived lymphocytes are isolated from the MLC mixture and administered intralesionally as an implant placed directly into the tumor of a patient desiring protection against recurrence of a tumor, for instance in the proximity of a surgically debulked tumor, or desiring treatment of an inoperable tumor. Alternatively, the alloactivated

donor lymphocytes can be administered peripherally in treatment of a tumor, such as at a secondary or metastatic tumor site. In another embodiment, the allogenic activated lymphocytes are co-cultivated with patient-derived lymphocytes, and the mixture of MLCs is implanted or introduced
5 peripherally as a mixture.

A BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 shows survival times of glioblastoma patients treated with intratumor implant of MLC cells.

5

FIGURE 2 is a graph showing MRI scans indicating the reduction in tumor size in nine glioblastoma patients, two patients, one receiving a single implant dose of 4×10^9 MLCs and one receiving a single implant dose of 6×10^9 MLCs, have continued to date to show a progressive reduction in tumor mass

10 over periods of 58 and 74 weeks, respectively.

A DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a method for treating tumors in mammals using an allograft of donor lymphocytes that have been co-cultured *in vitro* in a mixed lymphocyte (MLC) reaction with the patient's mononuclear cells, preferably from peripheral blood, to activate the donor lymphocytes against antigens associated with the patient's tumor. It is believed that a graft v. host response and a host v. graft response to the allograft induce a powerful immunologic reaction in the tumor site that results in cytokine production and tissue destruction. During this reaction the host lymphoid cells identify both the graft lymphoid cells and tissue as foreign and both are inhibited or destroyed by local and systemic immune response. This host reaction occurs not only at the implant site of the graft lymphoid cells, but may also occur peripherally, such as at a secondary or metastatic tumor site. In addition, the implanted, donor lymphocytes immunologically attack the tumor in a graft v. host response. Thus, a strong immunologic reaction in the center of the tumor produces an environment that overcomes the cytokine inhibitors that may be present at the tumor site. The increase in immunologic reaction will inhibit growth, decrease size and/or eradicate established tumors or developing metastases, as well as inhibit recurrence at the site of surgically debulked tumors. Thus, the present invention provides a method for potentiating the systemic immune response to the patient's tumor by introducing into the patient a viable preparation of the alloactivated cells described herein.

25

Numerous studies have shown that the *in vitro* environment in a mixed lymphocyte reaction facilitates an active primary immune response to allogeneic and tumor associated antigens (M. Gately, *et al.*, *JNCI* 69:1245, 1982; S. Lee, *et al.*, *J. Experimental Medicine* 147:912, 1978; J. Zarling, *et al.*, *Nature* 262:691, 1976). In the present invention, the MLC reaction

30

occurs within the patient's own tumor tissue, thereby stimulating the patient to respond against its own tumor. Preferably the co-culture is performed, generally from 1 to 5 days. Since release of key cytokines takes place during the early stages of the co-culture, it is preferred that the co-cultured cells be

5 implanted in the early stages of the co-culture reaction, usually during the first 48 hours of the reaction when cytokine levels are normally highest. This method results in release of key cytokines directly into the tumor tissue so that an environment is created that is conducive to antigen recognition and development of cell mediated immunity directed against the antigen.

10 Standard techniques, such as those based on immunoassay, can be used to measure the level of various cytokines, including TNF, LT and gamma interferon, present in the MLC culture supernatants. Levels of TNF and LT vary from 50 to 150 units of biologic activity/ml or 500 to 3500 pg/ml of supernatant. As a result of the *in vitro* co-culture, healthy allogenic lymphoid

15 cells from a donor are specifically activated against the patient's tumor associated antigens.

The implanted cells are generally allowed to come into direct contact with tumor cells. Thus, as used herein the term "implant" and "implanted" means

20 that the cells are placed into the patient's body as a group, optionally within a thickening matrix of coagulated serum or other suspension or gel-like substance.

Examples of tumors that can be treated by the method of this invention

25 include the following:

Brain tumors, such as astrocytoma, oligodendroglioma, ependymoma, medulloblastomas, and PNET (Primitive Neural Ectodermal Tumor);

Pancreatic tumors, such as pancreatic ductal adenocarcinomas.

Lung tumors, such as small and large cell adenocarcinomas, squamous cell carcinoma, and bronchoalveolarcarcinoma;

Colon tumors, such as epithelial adenocarcinoma, and liver metastases of these tumors;

5 Liver tumors, such as hepatoma, and cholangiocarcinoma;

Breast tumors, such as ductal and lobular adenocarcinoma;

Gynecologic tumors, such as squamous and adenocarcinoma of the uterine cervix, and uterine and ovarian epithelial adenocarcinoma;

Prostate tumors, such as prostatic adenocarcinoma;

10 Bladder tumors, such as transitional, squamous cell carcinoma;

Tumors of the RES System, such as B and T cell lymphoma (nodular and diffuse), plasmacytoma and acute and chronic leukemia;

Skin tumors, such as malignant melanoma; and

Soft tissue tumors, such as soft tissue sarcoma and leiomyosarcoma.

15

In one embodiment of the method of this invention, the tumor activated allogenic donor lymphocytes are implanted at the tumor site, optionally together with the patient's PBMCs from the co-culture, to supplement and enhance the patient's immune system attack upon autologous tumor cells.

20 For instance, in one embodiment of the invention alloactivated human donor lymphocytes obtained by co-cultivation with patient-derived PBMCs or cell lysates of PBMCs are administered intralesionally in a human, **for instance** as an implant placed directly into the brain of a patient desiring protection against recurrence of a glioblastoma tumor. The tumor activated allogenic

25 donor lymphocytes can be implanted in the proximity of a surgically debulked tumor, or a tumor treated by irradiation, chemotherapy, or other appropriate techniques. In another preferred embodiment, the allogenic activated lymphocytes are co-cultivated with patient-derived lymphocytes and the mixture of co-cultured allogenic and autologous cells, known herein as

30 "the MLCs," is implanted intralesionally.

The typical implant comprises a therapeutic amount of the allostimulated donor lymphocytes. As used herein the term "a therapeutic amount" means a sufficient quantity of MLCs or donor allostimulated lymphoid cells obtained from the mixed lymphocyte culture to inhibit growth, decrease the size of
5 and/or eradicate established tumors, and/or prevent tumor recurrence at the site of a tumor that has been surgically debulked or treated with chemotherapy or irradiation. More generally, a therapeutic amount may vary with the potency of each batch of alloactivated donor cells; the amount required for the desired therapeutic or other effect, the mode of
10 administration, *i.e.*, whether by direct implant into a tumor or body cavity or by peripheral administration, such as intravenously; and the rate of elimination or breakdown of the MLCs by the body once implanted or administered. In accordance with conventional prudent formulating practices, a dosage near the lower end of the useful range may be employed initially
15 and the dosage increased or decreased as indicated from the observed response, as in the routine procedure of the physician. In general, however, a unit dosage for direct implant comprises from about 2×10^9 to about 6×10^9 MLCs. For instance, it has been discovered that in the treatment of brain tumors, the upper limit of cells that can be implanted is about 6×10^9 .
20 Alternatively, a unit dosage for peripheral administration usually comprises from about 2×10^9 to about 2×10^{10} MLCs.

The invention further comprises a sterile vial or other container holding a composition comprising a unit dosage of the MLCs. Typically, the vial or
25 container will bear a label sets forth information concerning the pharmaceutical use of the composition in treating a tumor in a human, such as FDA approval for use of the composition in the treatment of a human having one or more of the tumors against which the method of treatment of the invention is effective as described herein.

Although any known method of obtaining PBMCs from a donor can be used, it is preferred to obtain approximately 150 to 300 ml of leukapheresis suspension containing the donor PBMCs, utilizing techniques of leukapheresis that are well known in the art for supportive apheresis according to the instructions of the manufacturer of the leukapheresis equipment. For instance, leukapheresis can be performed using a Cobe 2997, Cobe Spectra® (Lakewood, CO), Fenwall CS 3000 (Deerfield, IL), or Haemonetics (Braintree, MA) blood cell separator. Generally, a flow rate of 40 to 50 ml/min for 2 to 4 hours with lymphocyte yield of $2-4 \times 10^9$ can be used to process a total donor blood volume of 7,000 to 12,000 ml to yield 200 to 250 ml of leukapheresis suspension having less than 1 ml of red cells. For example, if a Cobe 2997 blood cell separator is used, the centrifuge rate is generally about $5 \times g$, the flow rate is up to about 45 ml/min and the collection rate is no more than or equal to 2.5 ml/min. One skilled in the art will appreciate that the yield of lymphocytes will vary with the donor and the leukapheresis machine used. For instance, if the donor pre-absolute lymphocyte counts are in the 0.6×10^9 to 1.0×10^9 level, as little as 150 ml of donor leukapheresis suspension can be drawn.

If desired, the donor cells can be contacted with a stimulatory cytokine such as IL-2 to trigger activation of the cells to the patient-derived antigen during co-culture and to further stimulate lymphocyte proliferation.

Donor PBMCs are obtained from the donor blood fraction, exercising care to avoid rupture of mononuclear cells, for instance by centrifuging the blood fraction containing the mononuclear cells through a cell separation medium such as Histopaque® 1.077 at $350 \times g$ for 7 to 10 minutes. Those of skill in the art will know of other PBMC cell separation techniques that can readily be utilized. Donor blood is typically pre-screened 3-7 days prior to surgery for HIV, Hepatitis A, B, and C, and VDRL.

Sufficient anticoagulant, such as 2% citrate, is added to the donor and patient blood or blood fraction to prevent coagulation upon withdrawal. Alternative anticoagulants and mixtures thereof known to one of skill in the art can also be used, such as anti-coagulant citrate dextrose formula A (ACDA) 15
5 ml/citrate 100 ml; anti-coagulant citrate dextrose formula B (ACDB) 25
ml/citrate 100 ml; or citrate phosphate dextrose (CPD) 14 ml/citrate 100 ml.

Typically, whole blood is withdrawn from the patient to be treated according to the invention using methods known in the art, such as vena puncture.
10 PBMCs are isolated from patient whole blood, usually by centrifugation through a cell separation medium, such as Histopaque® 1.077 (Sigma, St. Louis, MO), and are thoroughly washed to free the cells of the clotting factor in the patient's blood.

15 Samples of both the donor's and patient's blood or blood fraction should be thoroughly tested to ensure sterility before co-culturing of the cells. Typical of the tests for sterility of blood components that can be conducted by one of skill in the art, are those using such growth media as thioglycollate broth, tryptic soy broth and Roswell Park Memorial Institute Tissue Culture Medium
20 (RPMI) with 10% heat-inactivated fetal bovine serum (FBS)(RPMI - 10%) and 1% L-glutamine with no added antibiotics. Sterility tests utilizing the culture of blood cells in such growth media are illustrated in Example 1 of this application. Alternative sterility tests will be known to those of skill in the art.

25

Before mixed lymphocyte culture can be performed, typically the PBMCs isolated from whole blood are analyzed to determine the number of live cells per unit volume. This can be performed, for example, by using a stain that differentiates between living and dead cells and counting the cells in a
30 Neubauer chamber. Typical stains for this use are trypan blue and Eosin Y

dyes, both of which can be used in wet preparations. Alternative stains will be known to those of skill in the art. Generally, the concentration of live cells is standardized by diluting the preparation of PBMCs to achieve a predetermined concentration of live cells per unit volume. Although one
5 skilled in the art may select a number somewhat higher or lower, it is generally preferred that the number of live cells be fixed at a concentration of about 10^7 cells/ml for the purposes of conducting the mixed lymphocyte culture. The donor cells are typically cultured at a ratio of 10:1 to 20:1 compared to patient cells.

10

Standard techniques for conducting mixed lymphocyte culture using mammalian (*i.e.*, human) cells are well known in the art and are illustrated in Example 1 of this application. See, for instance, *Current Protocols in Immunology*, Ed. J. E. Coligan, *et al.*, John Wiley & Sons, Inc., 1994, Sec.
15 7.10 and M. Gately, *et al.*, *supra*; S. Lee, *et al.*, *supra*; and J. Zarling, *et al.*, *supra*, which are incorporated herein by reference in their entirety. To block response of the patient stimulator cells to the donor responder cells (back stimulation), it is preferred that the patient cells be irradiated or treated with a DNA binding agent, such as mitomycin C, before mixture of the cells
20 during co-culture to reduce or eliminate patient cell proliferative potential, as is well known in the art. In the present invention, donor lymphocytic cells are typically co-cultured in a short term mixed lymphocyte culture with the patient's PBMCs for a period of at least 48 hours, and preferably from 1 to 5 days. As a result of the *in vitro* co-culture, healthy lymphoid cells from the
25 unrelated donor are specifically activated against the patient's tumor associated antigens.

Preparations for parenteral or intravenous administration are typically contained in a "physiologically compatible carrier." Since the cells utilized
30 in the practice of this invention are live, a physiologically compatible carrier

is one that does not impair viability of the cells, *i.e.*, is hypotonic and at physiological pH. Such carriers include sterile aqueous salt solutions, suspensions and emulsions, including saline and buffered media, Ringer's dextrose, dextrose and sodium chloride, and lactated Ringer's solution.

- 5 Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers, such as those based on Ringer's dextrose, and the like. For administration by non-intravenous routes, the carrier can be in the form of clotted plasma, preferably the patient's clotted plasma. Alternatively the carrier can be a plasma-free, physiologically compatible, biodegradable solid
10 or semi-solid, such as a gel, suspension or water soluble jelly. Acacia, methylcellulose and other cellulose derivatives, sodium alginate and tragacanth suspensions or gels are suitable for use as carriers in the practice of this invention, for example, sodium carboxymethylcellulose 2.5%, tragacanth 1.25% and guar gum 0.5%.

15

- In a preferred method of implant at a tumor site, the co-cultured lymphocytes are collected from the co-culture supernatant (after at least 48 hours of co-culture) by centrifugation at the time of surgery. The collected cells are washed two times with injectable saline and re-suspended in platelet free,
20 decalcified plasma obtained from the patient the previous day. The cells in plasma are transported to surgery, the plasma is re-calcified by the addition of calcium, preferably in the form of calcium gluconate so that the plasma clots, embedding the cells in an autologous plasma clot. The patient's tumor is then surgically debulked, and the clot is aseptically minced and implanted
25 into the tumor site.

- The following examples illustrate the manner in which the invention can be practiced. It is understood, however, that the examples are for the purpose of illustration and the invention is not to be regarded as limited to any of the
30 specific materials or conditions therein.

EXAMPLE 1*A. Mixed Lymphocyte Cell (MLC) Culture Procedure.*5 1. Collection of responder PBMC from unrelated donor.

Peripheral blood mononuclear cells (PBMCs) were collected by leukapheresis from normal healthy donors unrelated to the patient. Donors were pre-screened to test for complete blood count (CBC) with differential, Hepatitis
10 A, B, and C, VDRL, and HIV-I.

Approximately 150 to 300 ml of leukapheresis suspension containing PBMC was collected from each donor, using standard blood donation procedures for supportive apheresis according to the manufacturers' instructions. The
15 leukapheresis was performed using a Fenwall CS 3000 (Deerfield, IL) blood cell separator. A flow rate of 40 to 50 ml/min for 2 to 4 hours with lymphocyte yield of $2-4 \times 10^9$ processed a total donor blood volume of 7,000 to 12,000 ml to yield 200 to 250 ml of leukapheresis suspension having less than 1 ml of red cells. If a Cobe 2997 blood cell separator was used, the
20 centrifuge rate was $5 \times g$, the flow rate was up to 45 ml/min, and the collection rate was no more than or equal to 2.5 ml/min.

However, if donor pre-absolute lymphocyte counts were in the 0.6×10^9 to 1.0×10^9 range, as little as 150 ml of leukapheresis product was drawn.
25 Hematocrit for the final product was 3.5%. At least one total blood volume was processed for 80% efficiency of lymphocyte collection.

The anticoagulant used was either 2% citrate or a citrate/anticoagulant ratio of ACDA - 15 ml/citrate - 100 ml; ACDB - 25 ml/citrate - 100 ml; or CPD
30 - 14 ml/ citrate - 100 ml. To obtain the utmost product purity, the actual and

final product from the cell separator was transported as a pure concentrate of cells in autologous plasma. The cells were not washed, and no albumin was added.

5 2. Preparation of donor cells.

The leukapheresis product was transported to the MC Oncology Research Laboratory for the production of allogeneic mixed lymphocyte cells (MLCs) for immunotherapy.

10

Cells were drained from the leukapheresis pack into two or three 250 ml centrifuge tubes, removing and setting aside 3 ml for sterility tests to be done during centrifugation. Cell concentrate was diluted with phosphate buffered saline (PBS) and centrifuged for 7 minutes at 2,000 rpm. Centrifugation was
15 repeated twice for a total of three times to wash the cells free of the clotting factor in the donor's serum.

Three 1 ml aliquots from the 3 ml removed from the leukocyte suspension were placed into sterile capped tubes for sterility testing. The first 1 ml
20 aliquot was added to thioglycollate medium (Difco, Detroit, MI) (30-35°C, 48 hr.); a second 1 ml was added to tryptic soy broth (Difco, Detroit, MI) (25-30°C, 48 hr.); and the third 1 ml was added to RPMI 1640 (GIBCO, Gaithersburg, MD) with 10% heat-inactivated FBS (RPMI-10%) and 1% L-glutamine, but without antibiotics.

25

Cells were spin washed twice at 150g for 10 minutes in PBS to remove platelets. The supernatant was very carefully discarded as cells were in a slurry and not a pellet. Cells were resuspended in AIM V (GIBCO, Gaithersburg, MD) supplemented with 2% heat inactivated FBS (2% AIM V)
30 to 420 ml, and placed into a T-175 cm² flask.

Patient or donor blood was diluted 1:1 with sterile saline. For cell separation, 35 ml of cell suspension was carefully layered onto 15 ml Histopaque® 1.077 suspension medium (Sigma, St. Louis, MO) in each 50 ml tube and centrifuged at 250g for 45 minutes. Centrifugation was started slowly and gradually increased to full speed. After centrifugation, the interface containing mononuclear cells between the Histopaque® suspension medium and the plasma layer was carefully collected with a 25 ml sterile pipet, deposited into clean 50 ml centrifuge tubes, diluted with 2% AIM V Media 1:1, and centrifuged at 550g for 7 to 10 minutes to form a cell pellet. Cells remained a minimum of time in the Histopaque® suspension medium, because it is toxic to the cells.

The supernatant was discarded, the pellet was resuspended in 2% AIM V and divided into two 50 ml centrifuge tubes to a total volume 40 ml, and centrifuged at 550g for 5 minutes. After washing, the supernatant was discarded. The washing step was repeated twice for a total of three times. After the last wash, cells in each tube were resuspended in 50 ml of 2% AIM V. Aliquots of 1 ml of the resuspended cells were diluted to a ratio of 1:10 in 2% AIM V per tube, then further diluted 1:1 in Trypan Blue (Sigma, St. Louis, MO) to distinguish dead from live cells, and the live cells were counted in a hemocytometer. Cells were set at 2×10^6 /ml with 2% AIM V.

3. Collection of stimulator PBMC from tumor patients.

From 200 to 400 ml of peripheral blood cells were drawn from each glioblastoma patient by vena puncture and placed into 250 ml centrifuge tubes, removing and setting aside 3 ml for sterility tests to be done during spinning. Blood cells in the centrifuge tubes were diluted with saline and centrifuged for 7 minutes at 550g. Centrifugation was repeated twice for a

total of three times to wash the cells free of the clotting factor in the patient's serum. Sterility testing was conducted as described above.

Cells were washed twice by centrifugation at 150g for 10 minutes in saline
5 to remove platelets, the supernatant was very carefully discarded, and 420 ml of cells were resuspended in a T-175 cm² flask in saline.

15 ml of Histopaque® 1.077 cell separation medium was added to twelve 50 ml centrifuge tubes, and 35 ml of cells suspended in saline were layered onto
10 the Histopaque® 1.077 in each 50 ml tube. The cell suspensions were spun at 250g for 45 minutes, starting centrifugation slowly and gradually increasing speed.

After centrifugation, the mononuclear cells at the interface between the
15 Histopaque® cell separation medium and the plasma layer were carefully collected with a 25 ml sterile pipet into 2 sterile 250 ml centrifuge tubes and diluted with 2% AIM-V to a final volume of 250 ml. The diluted mononuclear cells were centrifuged at 550g for 7 to 10 minutes. For washing, the supernatant was discarded, then the cell pellet was re-suspended
20 with 2% AIM V and centrifuged at 550g for 5 minutes. The washing step was repeated for a total of three times.

After the last washing step, cells were re-suspended in 50 ml of 2% AIM V, 1 ml of the cell suspension was diluted 1:10 in 2% AIM-V per tube, and the
25 number of viable cells was determined by enumeration in a 1:1 in Trypan Blue as described above.

The above-described procedures for collection of cells and proof of sterility have been approved by the State of California for GMP for sterility
30 and quality control.

4. Alloactivation of patient mononuclear cells (PBMC) with donor leukocytes.

- 5 The isolated patient PBMCs were re-suspended at 10^7 cells/ml in AIM-V, 50 μ g Mitomycin C (Bristol-Mayer Squibb, Princeton, NJ) were added per ml of patient cell suspension, and the suspension of PBMCs was incubated at 37 °C for one hour to block response of the stimulator cells to the responder cells (back stimulation). After one hour of incubation, the excess mitomycin
- 10 C was washed from the cells by alternate centrifugation (250g for 5 min), and the cells were resuspended in AIM-V. After mitomycin treatment of the patient's PBMCs, the cells were added at a 20:1 to 10:1 ratio to the donor culture (obtained as described above).
- 15 For co-culture, the donor and mitomycin C-treated patient PBMC suspension was placed in a sealed sterile Fenwal tissue culture system especially designed for culture of PBMC for reimplantation into patients. Cells were passed in sealed systems via Fenwal cell transfer units and pumps according to the manufacturers instructions, and cultured in a 37°C incubator for 48
- 20 hours.

5. Sterility testing of alloactivated cells.

- 25 Two days prior to implantation of the cell suspension, the following three sterility tests were performed. 10 ml sterile aliquots were removed from each tissue culture bag, placed into sterile capped 15 ml centrifuge tubes, and centrifuged for 10 minutes at 450g. In each tube, the pellet was resuspended in 3.0 ml of PBS. A 1 ml aliquot of the cell suspension was added to each
- 30 of three sterile capped tubes containing 2 ml of thioglycollate broth, tryptic

soy broth, or RPMI-10% and incubated for 48 hours. Each cell suspension was examined microscopically prior to implant to detect signs of microbial growth.

- 5 On the day of surgery, the cells were centrifuged out of their medium, washed two times with saline and re-suspended in platelet free, decalcified plasma obtained from the patient the previous day. The cells were transported to the operating room in plasma, then the plasma was re-calcified by the addition of calcium gluconate so that it clots just before implantation
10 into the tumor bed.

The day of surgery a drop of collected cell pellet was again examined for sterility under the microscope. Just prior to clotting, a 100 μ l aliquot of the cell suspension was added to 2 ml each of RPMI-10% without antibiotics,
15 thioglycollate and tryptic soy broth in a sterile capped tube. The samples were then incubated for four days after surgery, and a running log was kept of this last sterility test.

Cyto-implants produced by the above described method have received FDA
20 approval for use in human cancer patients (IND-BB 6288).

B. Intralesional implant of mixed lymphocyte cell culture.

25

A human Phase I trial was initiated to test the effects of intracranial injection of alloactivated allogeneic lymphoid cells in patients with recurrent glioblastoma. A total of 9 patients with recurrent Grade 3 and Grade 4 glioblastoma were entered into this trial. MLCs obtained from mixed
30 lymphocyte culture in Step 3 above were implanted intralesionally in the

debulked tumor at the time of surgery. Three patients were tested at each of three cell doses of 2×10^9 , 4×10^9 , and 6×10^9 of MLCs. The patients were followed clinically and magnetic resonance imaging (MRI) scans were performed at various monthly intervals on each patient following implantation
5 to monitor the progress of the disease.

All patients complained of headache and nausea for 1 to 2 months after surgery. Patients receiving the highest cell dose experienced the highest number of symptoms. Four of the five patients, including all those in the
10 highest dose and one each in the lower doses, showed a 50% or greater reduction in the area of tumor enhancement in MRI scans over a 1 to 3 month period after the implant. The results of MRI scans are shown in Figure 1, which measures survival time of glioblastoma patients treated with intratumor implants of MLC cells. Two patients, one receiving a single
15 implant dose of 4×10^9 MLCs and one receiving a single implant dose of 6×10^9 MLCs, have continued to date to show a progressive reduction in tumor mass over periods of 58 and 74 weeks, respectively. These two patients also have shown a marked improvement in Karnofsky score. The results of MRI scans are shown in Figure 2, which measures the reduction of tumor volume
20 as a function of time since implant of MLCs.

EXAMPLE 2

25 Phase I studies have been conducted in two patients with systemic melanoma. Mixed cultures of patient and donor lymphocytes were prepared as described in Example 1. The patients were treated by intralesional injection of 2×10^9 MLC cells into cutaneous tumors. Neither patient exhibited negative effects. The first patient experienced necrosis and destruction of the injected
30 cutaneous tumor and inflammation at a distant metastasis. The second patient

showed inflammation at the injected tumor, but no necrosis at the present dosage level. Treatment of these patients is ongoing.

EXAMPLE 3

5

A human Phase I clinical study has been conducted to test the effect of intratumor implant of MLC cells in patients with pancreatic cancer. Four patients with incurable, untreatable pancreatic cancer received intralesional implant of 4×10^9 MLC cells. Little or no toxicity was found. A reduction
10 in tumor mass occurred and extension of life span has been observed. Three patients have experienced a greater than 50 percent reduction in tumor mass. In patient No. 3, the serum levels of CA-19-9, a marker for this tumor, declined from a high of 206 to 86 over a two month period. Seven months after treatment, two of the treated patients have returned to their jobs, have
15 no clinical symptoms of tumor, and are living a totally normal life.

The foregoing description of the invention is exemplary for purposes of illustration and explanation. It should be understood that various modifications can be made without departing from the spirit and scope of the
20 invention. Accordingly, the following claims are intended to be interpreted to embrace all such modifications.

CLAIMS

1. A method of treating a mammalian patient having a tumor comprising administering to the patient a therapeutically effective amount of
5 allogenic donor lymphocytes that have been co-cultured in the presence of patient-derived lymphocytes under conditions sufficient to alloactivate the donor lymphocytes.
2. The method of claim 1 wherein proliferation of the patient derived
10 lymphocytes is blocked.
3. The method of claim 2 wherein the proliferation is blocked with mitomycin C.
- 15 4. The method of claim 1 wherein the tumor is malignant.
5. The method of claim 4 wherein the tumor is neuronally derived.
6. The method of claim 5 wherein the patient is a human and the tumor
20 is a glioblastoma.
7. The method of claim 4 wherein the tumor is epidermally derived.
8. The method of claim 7 wherein the tumor is melanoma.
- 25 9. The method of claim 1 wherein the tumor is a primary tumor.
10. The method of claim 4 wherein the tumor is a metastasis.

11. The method of claim 1 wherein the allogenic donor lymphocytes are administered intralesionally.
12. The method of claim 11 wherein the therapeutically effective amount
5 of allogenic donor lymphocytes is in the range from about 2×10^9 to 6×10^9 .
13. The method of claim 1 wherein the patient is a human and the donor is a human.
10
14. The method of claim 13 wherein the patient is also treated by a method selected from the group consisting of surgical debulking of the tumor, chemotherapy and irradiation.
- 15 15. The method of claim 1 wherein the allogenic donor lymphocytes are administered peripherally.
16. The method of claim 15 wherein the therapeutically effective amount
20 of allogenic donor lymphocytes is in the range from about 2×10^9 to 2×10^{10} .
17. The method of claim 16 wherein the therapeutically effective amount of cultured lymphocytes is introduced embedded in clotted patient-derived plasma.
25
18. The method of claim 1 wherein the co-culture is in the presence of a lymphocyte proliferation agent.
19. The method of claim 18 wherein the lymphocyte proliferation agent
30 is IL-2.

20. A method of treating a human having a tumor comprising:
- a) collecting lymphocytes from a normal human donor;
 - b) collecting lymphocytes from a patient having a tumor;
 - c) contacting the donor lymphocytes and patient lymphocytes
5 under conditions sufficient to alloactivate the donor lymphocytes; and
 - d) introducing a therapeutically effective amount of cultured lymphocytes into the patient in the vicinity of the tumor.
- 10 21. The method of claim 20 wherein the contacting in step c) of the donor lymphocytes and the patient lymphocytes is for at least 48 hours.
22. The method of claim 21 wherein the proliferation is blocked with mitomycin C.
- 15 23. The method of claim 20 wherein the tumor is malignant.
24. The method of claim 23 wherein the tumor is neuronally derived.
- 20 25. The method of claim 24 wherein the tumor is a glioblastoma.
26. The method of claim 23 wherein the tumor is epidermally derived.
27. The method of claim 26 wherein the tumor is melanoma.
- 25 28. The method of claim 20 wherein the tumor is a primary tumor.
29. The method of claim 23 wherein the tumor is a metastasis.

30. The method of claim 20 wherein the allogenic donor human lymphocytes are administered intralesionally.
31. The method of claim 30 wherein the therapeutically effective amount
5 of allogenic donor human lymphocytes is in the range from about 2×10^9 to 6×10^9 .
32. The method of claim 20 wherein the patient is also treated by a
10 method selected from the group consisting of surgical debulking of the tumor, chemotherapy and irradiation.
33. The method of claim 20 wherein the allogenic donor human lymphocytes are administered peripherally.
- 15 34. The method of claim 33 wherein the therapeutically effective amount of allogenic donor human lymphocytes is in the range from about 2×10^9 to 2×10^{10} .
35. The method of claim 20 wherein the therapeutically effective amount
20 of cultured lymphocytes is introduced embedded in clotted patient-derived plasma.
36. The method of claim 20 wherein the co-culture is in the presence of a lymphocyte proliferation agent.
- 25 37. The method of claim 36 wherein the lymphocyte proliferation agent is IL-2.

38. A therapeutic composition for inhibiting tumor cell proliferation in a mammalian patient comprising a combination of cells in a substantially plasma-free physiologically compatible carrier comprising a first lymphocyte population derived from a human patient having a tumor and a second lymphocyte population derived from a normal allogenic donor, wherein the donor lymphocytes are alloactivated to the to the tumor.
39. The composition of claim 38 wherein the carrier is a gel.
40. The composition of claim 38 wherein the composition is substantially sterile.
41. The composition of claim 38 wherein the composition comprises a unit dosage of the cells.
42. The composition of claim 41 wherein the unit dosage is from about 2×10^9 to about 6×10^9 of the cells.
43. The composition of claim 41 wherein the unit dosage is from about 2×10^9 to about 2×10^{10} of the cells.

44. An article of manufacture for use in treating tumors in a mammalian host comprising:

(a) a sterile vial; and

5 (b) a composition contained within the vial comprising a combination of cells in a physiologically compatible carrier comprising a first lymphocyte population derived from a patient having a tumor and a second lymphocyte population derived from a normal allogenic donor, wherein the donor lymphocytes are alloactivated to the tumor.

10


45. The article of claim 44 wherein the host is a human.

46. The article of claim 44 further comprising a label on the sterile vial,
15 wherein the label sets forth information concerning the pharmaceutical use of the composition in treating a tumor in a human.

47. The article of claim 44 wherein the host is a human.


1/2

FIG. 1



2/2

FIG. 2



WEEKS, 0 POINT = DATE OF IMPLANT (LAST DATA POINT FOR ALL PATIENTS
REPRESENT TOTAL WEEKS SURVIVAL SINCE LAST MRI)

TUMOR VOLUME CALCULATED FROM AREAS OF
ENHANCEMENT (CM³), MEASURED BY DR. JACQUES

INTERNATIONAL SEARCH REPORT

International application No.

PC 96/03621

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 5/00; A61K 45/05, 48/00, 31/40

US CL : 435/240.1; 424/85.2, 93.21; 514/410

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/240.1; 424/85.2, 93.21; 514/410

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS cytotoxic and Dupont; Dupont, 7/in and lymphocyte?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4,677,056 A (B. DUPONT ET AL.) 30 June 1987 (30.06.87) columns 7 and 8.	1-47



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

18 JUNE 1996

Date of mailing of the international search report

02 JUL 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOHN W. ROLLINS

Telephone No. (703) 308-0196

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999

1000